

California School Employee: Tuberculosis (TB) Risk Assessment

Job-related requirement for child care, pre-K, K-12, and Volunteers

- Use of this questionnaire is required by California Education Code sections 49406 and 87408.6, and Health and Safety Code sections 1597.055 and 121525-121555.
- The purpose of this tool is to identify **adults** with infectious tuberculosis (TB) to prevent them from spreading disease.
- **Do not repeat testing** unless there are **new risk factors since the last negative test**.
- Do not treat for latent TB infection (LTBI) until active TB disease has been excluded:
For individuals with signs or symptoms of TB disease or abnormal chest x-ray consistent with TB disease, evaluate for active TB disease with a chest x-ray, symptom screen, and if indicated, sputum AFB smears, cultures and nucleic acid amplification testing. A negative tuberculin skin test (TST) or interferon gamma release assay (IGRA) does not rule out active TB disease.

The law requires that a health care provider administer this risk assessment. A health care provider, as defined for this purpose, is any organization, facility, institution or person licensed, certified or otherwise authorized or permitted by state law to deliver or furnish health services. Any person administering this risk assessment is to have training in the purpose and significance of the risk assessment and Certificate of Completion.

Name of Employee/Volunteer Assessed for TB Risk Factors: _____

Classified Certificated

Employee PIN: _____

Date of assessment/examination: _____ mo/_____ day/_____ year Date of Birth: _____ mo/_____ day/_____ year

History of Tuberculosis Infection or Disease (Check appropriate box below)

- Yes** If there is a documented history of positive TB test or TB disease, then a symptom review and chest x-ray (if none performed in previous 6 months) should be performed at initial hire by a physician, physician assistant, or nurse practitioner. If the x-ray does not have evidence of TB, the person is no longer required to submit to a TB risk assessment or repeat chest x-rays.
- No** (Assess for Risk Factors for Tuberculosis using information below)

TB testing is recommended if any of the 3 boxes below are checked.

- One or more signs and symptoms of TB disease**
- TB symptoms include prolonged cough, coughing up blood, fever, night sweats, weight loss, or excessive fatigue.
- Birth, travel, or residence** in a country with an elevated TB rate for at least 1 month
- Includes countries other than the United States, Canada, Australia, New Zealand, or Western and North European countries.
 - Interferon gamma release assay (IGRA) is preferred over tuberculin skin test (TST) for non-US-born persons.
- Close contact** to someone with infectious TB disease at any time.

The above named patient has submitted to a tuberculosis risk assessment. The patient does not have risk factors, or if tuberculosis risk factors were identified, the patient has been examined and determined to be free of infectious tuberculosis.

Signature of Health Care Provider completing the risk assessment and/or examination

Please print, place label or stamp with Health Care Provider Name and Address (include Number, Street, City, State, and Zip Code):